

CHEMICAL AND DYNAMICAL EVOLUTION IN THE SOLAR NEIGHBOURHOOD

O. Bienaymé¹, T. Mishenina², V. Kovtyukh², C. Soubiran³

¹Strasbourg Observatory, France, *bienayme@astro.u-strasbg.fr*

²Odessa Observatory, Ukraine

³Bordeaux Observatory, France

ABSTRACT. We measure the surface mass density of the galactic disk in the solar neighbourhood using two samples of red clump stars, we also determine the thin and thick disk chemical properties. High resolution spectral data of red clump stars towards the NGP have been obtained with the spectrograph Elodie at OHP for Tycho-2 selected stars. Nearby Hipparcos red clump stars were also observed with the Elodie spectrograph.

Keywords: Stars: abundances and kinematics – Galaxy: stellar populations.

1. Introduction

This paper is the extension of the previous works published by (Soubiran et al., 2003) and (Siebert et al., 2003) to probe the properties and the distribution of red clump stars within 100 pc from the sun and at larger distances towards the north galactic pole.

2. The survey

To study the kinematics properties of the galactic disk and to determine the vertical force perpendicular to the galactic plane, we measure the vertical spatial distribution and the vertical velocity distribution of a test stellar population. As far as possible, this test population must be homogeneous and unbiased with selection criteria independent of velocities and distances. It must also be in a stationary state. For this purpose, we use a sample of red clump stars, extending to larger distances from the galactic plane the NGP sample previously analysed in (Soubiran et al., 2003; Siebert, 2003).

This sample is built from a preliminary list of red clump candidates, a list based on the Tycho-2 star catalogue (Hög et al., 2000) for which we select all stars in a restricted range of colour ($B - V$ within 0.9 and 1.1) in two fields in directions close to the North Galactic Pole. This preliminary list is dominated by

red clump stars: our high resolution spectroscopic observations allow to determine the absolute magnitudes and to confirm and separate the red clump stars from other stars. We have also improved the local counterpart sample of red giants by measuring and determining the radial velocities and metallicities for a complete set of 203 Hipparcos red clump stars.

2.1. The Hipparcos red clump stars

We select the 203 nearby red clump Hipparcos stars with a parallax larger than 10 mas, an equatorial latitude larger than -20 deg, a $B_J - V_J$ Johnson colour within 0.9 – 1.1 (the Johnson magnitudes being determined from the Tycho-2 B_T and V_T magnitudes applying Eq. 1.3.20 from (ESA SP-1200):

$$\begin{aligned} V &= V_T - 0.090 (B_T - V_T) \\ B - V &= 0.850 (B_T - V_T) , \end{aligned} \quad (1)$$

and an absolute magnitude M_{V_j} within 0. – 1.3. Most of these stars were observed with the echelle spectrograph Elodie in February 2003, October 2003 and February 2004 at the Observatoire de Haute Provence (France). The signal to noise ratios range from 150 to 200. We have measured and determined their radial velocity, the atmospheric parameters (T_{eff} , $\log g$, microturbulent velocities V_t , $[\text{Fe}/\text{H}]$), and abundances of some elements. The abundances of Fe, Ca, Si, and Ni have been determined from equivalent widths under LTE approximation whereas abundances of Mg have been determined under NLTE approximation using equivalent widths of 4 lines and profiles of 5 lines.

2.2. NGP K giants

The distant K giant sample is drawn from the Tycho-2 star catalogue. We have applied the same criteria given in (Soubiran et al., 2003) to build the list of red clump candidates, just extending the limiting apparent magnitudes to fainter stars. In summary, we have extracted from the Tycho-2 catalogue, stars in two fields

close to the north galactic pole. The first field is circular with a 10 deg radius and is centered towards the galactic direction $b = +90$ deg. The second field is also circular (radius 15 deg) and centered on the galactic direction ($l = 35.5$ deg, $b = +80$ deg) (we have removed stars close to the Coma open cluster direction, $l = 221$ deg, $b = 84$ deg, inside a 4.5 deg radius circle). The total area effectively covered by our samples is 720 square degrees.

From these two fields, we select stars with a $B_J - V_J$ colour within 0.9 – 1.1. For the first field we keep stars with an apparent magnitude V_J within 7 and 10.6 and for the second one with V_J within 7. to 9.5. About 500 candidates have been observed with the echelle spectrograph Elodie at OHP with a median S/N ratio of 20. This low S/N is sufficient to obtain the stellar parameters, T_{eff} , gravity and $[\text{Fe}/\text{H}]$ metallicity, and also the absolute magnitude by comparison and fitting to high resolution spectra within the TGMET library (Katz et al., 1998). The determined absolute magnitudes are then used to identify the real red clump giants and remove the dwarfs and subgiants.

3. Discussion

Nearly 700 Tycho-2 stars have been observed in the solar neighbourhood at distances smaller than 100 pc or in a 720 square degree field in the direction of the North Galactic Pole. Absolute magnitudes, effective temperatures, gravities and metallicities have been estimated, as well as distances and 3D velocities. Abundances of Fe, Si and Ni have been determined from equivalent widths under LTE approximation, whereas abundances of Mg have been determined under NLTE approximation. Most of these stars are clump giants and span typical distances from 0 pc to 800 pc to the galactic mid-plane. This new sample, free of any kinematical and metallicity bias, is used to investigate the vertical distribution of disk stars.

The old thin disk and thick disk populations are deconvolved from the velocity-metallicity distribution of the sample and their parameters are determined. The thick disk is found to have a moderate rotational lag with respect to the Sun with a mean metallicity of $[\text{Fe}/\text{H}] = -0.48 \pm 0.05$ and a high local normalization of $15 \pm 7\%$. We also determine both the gravitational force law perpendicular to the Galactic plane and the total surface mass density and thickness of the Galactic disk. The surface mass density of the Galactic disk within 800 pc derived from this analysis is $\Sigma (|z| < 800 \text{ pc}) = 76 \text{ Msun pc}^{-2}$.

The thickness of the total disk mass distribution is dynamically measured for the first time and is found to be 390 pc in relative agreement with the old stellar disk scaleheight. All the dynamical evidences concerning the structure of the disk (its local volume density - i.e. the Oort limit-, its surface density and its thickness) are compatible with our current knowledge of the corresponding stellar disk properties. This result implies that the dark matter component of our Galaxy cannot be distributed in a flat or disk-like component but must be distributed in a round halo.

References

- Soubiran C., Bienaymé O., Siebert A.: 2003, *A&A*, **398**, 141.
 Siebert A., Bienaymé O., Soubiran C.: 2003, *A&A*, **399**, 531.
 Hög E., Fabricius C., Makarov V.V.: 2000, *A&A*, **355**, 27.
 The Hipparcos and Tycho Catalogues: 1997, *ESA SP-1200*, Vol. 1-17.
 Kovtyukh et al.: 2004, in preparation.
 Mishenina et al.: 2004, in preparation.
 Katz D., Soubiran C., Cayrel R. et al.: 1998, *A&A*, **338**, 151.